

## Letter to the Editor

# Comparison of Female and Male Interstitial Deletions in the Distal Xq

### To the Editor:

We reviewed female interstitial deletions in the distal Xq and compared them to those reported in males [Schmidt, 1996]. Most of the deletions were common to females and males, and they were scattered within Xq27 and proximal Xq. Six females had large deletions of 1–10 Mb which formed a contig covering ~13 Mb within Xq27.1→proximal Xq28 (Fig. 1). In 3 of these patients the deleted X chromosome was preferentially active, and the phenotype was abnormal (mental retardation in the patient of Schmidt et al. [1990]; mental retardation and Hunter syndrome in the patient of Clarke et al. [1992]; and mental retardation and myotubular myopathy in the patient of Dahl et al. [1995]). All three deletions occurred *de novo*. Our previous analysis of these deletions showed no abnormalities in the methylation and replication patterns of the region distal to the deletion, and in the corresponding area on the normal X chromosome [Schmidt et al., 1994]. Thus, there is no evidence that the skewed inactivation pattern in these cases resulted from the cell selection driven by anomalies of X inactivation.

Three other large deletions, and three overlapping small ones, were found in normal females. All of these were detected because of the affected male offspring to whom the deleted X chromosome had been passed. The boys were all mentally retarded, and one additionally had Hunter syndrome. In each of the carrier mothers the X inactivation pattern was random. This finding is remarkable, as the area covered by these deletions (~12 Mb) almost fully overlaps with that of the three deletions with skewed inactivation pattern. We conclude that deletions of Xq27→proximal Xq28 do not promote cell selection against the deleted X chromosome as active. The resulting inactivation pattern is established by chance. The above conclusion is supported by two other female deletions of Xq27 presented by Willard [1996]. This implies that phenotypically normal carriers may have affected daughters if the deleted X chromosome in the offspring is preferentially active.

It has been known that females with deletions of the terminal Xq are fertile [Skibsted et al., 1984; Tharapel et al., 1993]. Our comparison of female and male deletions confirms that breakpoints within Xq27.1→proximal Xq28 do not interfere with female fertility, since the deletions of Quan et al. [1995], Meijer et al. [1994], Gedeon et al. [1992, 1995], and Beck et al. [1992] were all passed to the offspring (Fig. 1). While it is too early to exclude premature ovarian failure in the carriers of these deletions, there is no clear evidence for it. In the case of Quan et al. [1995] the mother was 33, in the family of Meijer et al. [1994] 4 carriers had 3 or 2 children each (ages 19, 17, and 9 in one case), and in the family of Gedeon et al. [1992] 1 carrier had 3 children and another had 4, the youngest being 19.

In summary, mental status in females with interstitial deletions of terminal Xq is determined by the pattern of X inactivation, which is highly variable. In contrast, fertility may not be affected at all.

### REFERENCES

- Albright SG, Lachiewicz AM, Tarleton JC, Rao KW, Schwartz CE, Richie R, Tennison MB, Aylsworth AS (1994): Fragile X phenotype in a patient with a large *de novo* deletion in Xq27–q28. *Am J Med Genet* 51:294–297.
- Beck M, Steglich C, Zabel B, Dahl N, Schwinger E, Hopwood JJ (1992): Deletion of the Hunter gene and both DXS466 and DXS304 in a patient with mucopolysaccharidosis type II. *Am J Med Genet* 44:100–103.
- Clarke JTR, Wilson PJ, Morris CP, Hopwood JJ, Richards RI, Sutherland GR, Ray PN (1992): Characterization of a deletion at Xq27–q28 associated with unbalanced inactivation of the normal X chromosome. *Am J Hum Genet* 51:316–322.
- Dahl N, Hu LJ, Chery M, Fardeau M, Gilgenkrantz S, Nivelon-Chevallier A, Sidaner-Noisette I, Mugneret F, Gouyon J-B, Gal A, Kioschis P, D'Urso M, Mandel J-L (1995): Myotubular myopathy in a girl with a deletion at Xq27–q28 and unbalanced X inactivation assigns the MTM1 gene to a 600-kb region. *Am J Hum Genet* 56:1108–1115.
- De Graaff E, Rouillard P, Willems P, Smits APT, Rousseau F, Oostra BA (1995): Hotspot for deletions in the CGG repeat region of FMR1 in fragile X patients. *Hum Mol Genet* 4:45–49.
- Fuchs S, Sarde CO, Wedemann H, Schwinger E, Mandel J-L, Korn B, Poustka A, Yu S, Sutherland GR, Mulley JC (1992): Fragile X syndrome without CCG amplification has an FMR1 deletion. *Nat Genet* 1:341–344.
- Gedeon AK, Baker E, Robinson H, Partington MW, Gross B, Manca A, Korn B, Poustka A, Yu S, Sutherland GR, Mulley JC (1992): Fragile X syndrome without CCG amplification has an FMR1 deletion. *Nat Genet* 1:341–344.
- Gez J, Gedeon AK, Sutherland GR, Mulley JC (1996): Identification of the gene FMR2, associated with FRAXE mental retardation. *Nature Genet* 13:105–108.
- Gedeon AK, Baker E, Robinson H, Partington MW, Gross B, Manca A, Korn B, Poustka A, Yu S, Sutherland GR, Mulley JC (1992): Fragile X syndrome without CCG amplification has an FMR1 deletion. *Nat Genet* 1:341–344.

Received for publication October 24, 1995; revision received December 18, 1995.

Address reprint requests to Malgorzata Schmidt, Department of Genetics and Human Variation, La Trobe University, Melbourne 3083, Australia.




Fig. 1. Male (left) and female (right) interstitial deletions in the distal Xq. **A:** Map of X chromosome scaled in megabases. **B:** Deletions reported in males are represented by vertical lines with name of first author and approximate size in kb. **C:** Order of X chromosome markers [Willard et al., 1994]. Bold horizontal symbols indicate position of known genes involved in mental retardation. The position of SOX3 [Stevanovic et al., 1993] has been narrowed (Schmidt, unpublished) using an X chromosome with a duplication DXS152 → Xqter Schmidt et al., 1991. **D:** Deletions reported in females are grouped according to X-inactivation pattern (in French brackets). In each of those indicated as having nonrandom X-inactivation, the deleted X was active and the individual was mentally retarded. Name of first author and approximate size of deletion in kb are given. References: Fuchs < 1, Sarde et al., 1994; Gedeon 200, 100, Gedeon et al., 1995; Gedeon 2500, Gedeon et al., 1992; MTM1, Dahl et al., 1995; Schmidt 10,000, Schmidt et al., 1990; SOX3, Stevanovic et al., 1993; Tarleton ~5000, Albright et al., 1994; Wilson 1700, 800, Steen-Bondeson et al., 1992; Wohrle 250–Hirst 25, 0.6, Hirst et al., 1995 (deletions reported by de Graaff et al. [1995] are not included in Fig. 1, as they result from somatic instability of the expanded FMR1 gene).

- Gedeon AK, Keinanen M, Ades LC, Kaariainen H, Gez J, Baker E, Sutherland GR, Mulley JC (1995): Overlapping submicroscopic deletions in Xq28 in two unrelated boys with developmental disorders: Identification of a gene near FRAXE. *Am J Hum Genet* 56:907-914.
- Gu Y, Shen Y, Gibbs RA, Nelson DL (1996): Identification of FMR2, a novel gene associated with the FRAXE CCG repeat and CpG island. *Nature Genet* 13:109-113.
- Gu Y, Lugenbeel KA, Vockley JG, Grody WW, Nelson DL (1994): A de novo deletion in FMR1 in a patient with developmental delay. *Hum Mol Genet* 3:1705-1706.
- Hirst M, Grewal P, Flannery A, Slatter R, Maher E, Barton D, Fryns J-P, Davies K (1995): Two new cases of FMR1 deletion associated with mental impairment. *Am J Hum Genet* 56:67-74.
- Kenrick S, Levinson B, Taylor S, Shapiro A, Gitschier J (1992): Isolation and sequence of two genes associated with a CpG island 5' of the factor VIII gene. *Hum Mol Genet* 1:179-186.
- Meijer H, de Graaff E, Merckx ML, Jongbloed JE, de Die-Smulders EM, Englen JJM, Fryns J-P, Curfs PMG, Oostra BA (1994): A deletion of 1.6 kb proximal to the CGG repeat of the FMR1 gene causes the clinical phenotype of the fragile X syndrome. *Hum Mol Genet* 3:615-620.
- Palmieri G, Capra V, Romano G, D'Urso M, Johnson S, Schlessinger D, Morris P, Hopwood J, Di Natale P, Gatti R, Ballabio A (1992): The iduronate sulfatase gene: Isolation of a 1.2-Mb YAC contig spanning the entire gene and identification of heterogenous deletions in patients with Hunter syndrome. *Genomics* 12:52-57.
- Quan F, Zonana J, Gunter K, Peterson KL, Magenis RE, Popovich BW (1995): An atypical case of fragile X syndrome caused by a deletion that includes the FMR1 gene. *Am J Hum Genet* 56:1042-1051.
- Rosenthal A, Jouet M, Kenrick S (1992): Aberrant splicing of neural cell adhesion molecule L1 mRNA in a family with X-linked hydrocephalus. *Nat Genet* 2:107-112.
- Rousseau F, Vincent A, Rivella S, Heitz D, Triboli C, Maestrini E, Warren ST, Suthers GK, Goodfellow P, Mandel J-L, Toniolo D, Oberle I (1991): Four chromosomal breakpoints and four new probes mark out a 10-cM region encompassing the fragile X locus (FRAXA). *Am J Hum Genet* 48:108-116.
- Sarde C-O, Mosser J, Kioschis P, Kretz C, Vicaire S, Aubourg P, Poustka A, Mandel J-L (1994): Genomic organization of the adrenoleukodystrophy gene. *Genomics* 22:13-20.
- Schmidt M (1996): Letter to the Editor: Mapping of the loci for mental retardation syndromes in the distal Xq. *Am J Med Genet* 64:163-165.
- Schmidt M, Certoma A, Du Sart D, Kalitsis P, Leversha M, Fowler K, Sheffield L, Jack I, Danks D (1990): Unusual X chromosome inactivation in a mentally retarded girl with an interstitial deletion Xq27: Implications for the fragile X syndrome. *Hum Genet* 84:347-352.
- Schmidt M, Du Sart D, Kalitsis P, Leversha M, Dale S, Sheffield L, Toniolo D (1991): Duplications of the X chromosome in males: Evidence that most parts of the X chromosome can be active in two copies. *Hum Genet* 86:519-521.
- Schmidt M, Robertson A, Crawford M (1994): X inactivation in interstitial deletions of the fragile X region. *Am J Med Genet* 51:451-453.
- Skibsted L, Westh H, Niebuhr E (1984): X long-arm deletions. A review of non-mosaic cases studied with banding techniques. *Hum Genet* 67:1-5.
- Steen-Böndeson M-L, Dahl N, Tønnesen T, Kleijer WJ, Seidlitz G, Gustavson K-H, Wilson PJ, Morris CP, Hopwood JJ, Pettersson U (1992): Molecular analysis of patients with Hunter syndrome: Implication of a region prone to structural alterations within the IDS gene. *Hum Mol Genet* 1:195-198.
- Stevanovic M, Lovell-Badge R, Collignon J, Goodfellow PN (1993): SOX3 is an X-linked gene related to SRY. *Hum Mol Genet* 12:2013-2018.
- Tarleton JC, Richie R, Schwartz C, Rao K, Aylsworth AS, Lachiewicz A (1993): An extensive de novo deletion removing FMR1 in a patient with mental retardation and the fragile X syndrome phenotype. *Hum Mol Genet* 2:1973-1974.
- Tharapel AT, Anderson KP, Simpson JL, Martens PR, Wilroy RS, Llerena JC, Schwartz CE (1993): Deletion (X)(q26.1→q28) in a proband and her mother: Molecular characterization and phenotypic-karyotypic deductions. *Am J Hum Genet* 52:463-471.
- Trottier Y, Imbert G, Poustka A, Fryns J-P, Mandel J-L (1994): Male with typical fragile X phenotype is deleted for part of the FMR1 gene and for about 100 kb of upstream region. *Am J Med Genet* 51:454-457.
- Willard HF (1996): X chromosome inactivation and X-linked mental retardation. *Am J Med Genet* 64:21-26.
- Willard HF, Cremers F, Mandel J-L, Monaco AP, Nelson DL, Schlessinger D (1994): Report of the Fifth International Workshop on Human X Chromosome Mapping 1994. *Cytogenet Cell Genet* 67:296-328.
- Wilson PJ, Suthers GK, Callen DF, Baker E, Nelson PV, Cooper A, Wraight JE, Sutherland GR, Morris CP, Hopwood JJ (1991): Frequent deletions at Xq28 indicate genetic heterogeneity in Hunter syndrome. *Hum Genet* 86:505-508.
- Wohrle D, Kotzot D, Hirst MC, Manca A, Korn B, Schmidt A, Barbi G, Rott H-D, Poustka A, Davies KE, Steinbach P (1992): A microdeletion of less than 250 kb, including the proximal part of the FMR-1 gene and the fragile X site, in a male with the clinical phenotype of fragile-X syndrome. *Am J Hum Genet* 51:299-306.

**Malgorzata Schmidt**

Department of Genetics and Human Variation  
La Trobe University  
Melbourne, Australia